Substitute Form PT (Modified)

U.S. Department of Commerce Patent and Trademark Office

Attorney's Docket No. 16743-004001 Application No. 10/813,549

formation Disclosure Statement by Applicant (Use several sheets if necessary)

Applicant You-Di Liao

Filing Date

Group Art Unit

(37 CFR §1.98(b))

March 29, 2004 Unknown

	U.S. Patent Documents						
Examiner Initial	Desig. ID	Document Number	Publication Date	Patentee	Class	Subclass	Filing Date If Appropriate
wwm	AA	5,013,662	05/07/1991	Ben-Bassat, et al.	435	2/2	
	AB						
	AC	•					
	AD						
	AE						
	AF						
	AG						
	AH						
	ΑI	•				-	
	AJ						
	AK						

	Foreig	n Patent Doo	uments or Pu	blished Foreign	Patent A	Application	าร	
Examiner	Desig.	Document	Publication	Country or			Trans	lation
Initial	ID	Number	Date	Patent Office	Class	Subclass	Yes	No
	AL	•						
	AM							
	AN							
	AO	-					•	
	AP							

	Other Documents (include Author, Title, Date, and Place of Publication)			
Examiner Initial	Desig. ID	Document		
ann	AQ	Abe, A., et al. (2000) Acetylation at the N-terminus of actin strengthens weak interaction between actin and myosin. Biochem. Biophys. Res. Commun., 268:14-19.		
uum	AR	Adachi, K., et al. (2000) Expression of functional soluble human α-globin chains of hemoglobin in bacteria. Protein Expr. Purif., 20:37-44.		
WWW	AS	Belagaje, R.M., et al. (1997) Increased production of low molecular weight recombinant proteins in Escherichia coli. Protein Sci., 6:1953-1962.		

Examiner Signature	Date Considered
	Date Considered
MIMMAN IVUSOU	L December 2005
EXAMINER: Initials distribution considered. Draw line through citation if no	t in conformance and not considered. Include copy of this form with
next communication to applicant.	
	Substitute Disclosure Form (PTO-1449)

Substitute Form PTO-1449 U.S. Department of Commerce (Modified) Patent and Trademark Office		Attorney's Docket No. 16743-004001	Application No. 10/813,549	
Information Disclosure Statement by Applicant		Applicant You-Di Liao		
	neets if necessary)	Filing Date March 29, 2004	Group Art Unit Unknown	

	Other Documents (include Author, Title, Date, and Place of Publication)			
Examiner	Desig.	. Barrant		
Initial	ID	Document Ben-Bassat, A., et al. (1987) Processing of the initiation methionine from proteins:		
www	AT	properties of the <i>Escherichia coli</i> methionine aminopeptidase and its gene structure. J. Bacteriol., 169(2):751-757.		
	AU	Boix, E., et al. (1996) Role of the N terminus in RNase A homologues: differences in catalytic activity, ribonuclease inhibitor interaction and cytotoxicity. J. Mol. Biol., 257:992-1007.		
	AV	Busby, W.H., Jr., et al. (1987) An enzyme(s) that converts glutaminyl-peptides into pyroglutamyl-peptides. Presence in pituitary, brain, adrenal medulla, and lymphocytes. J. Biol. Chem., 262(18):8532-8536.		
	AW	Chang, S.Y., et al. (1989) Methionine aminopeptidase gene of Escherichia coli is essential for cell growth. J. Bacteriol., 171(7):4071-4072.		
	AX	Chen, S., et al. (2002) The specificity in vivo of two distinct methionine aminopeptidases in Saccharomyces cerevisiae. Arch. Biochem. Biophys., 398(1):87-93.		
	AY	Chiu, C.H., et al. (1999) Amino acid residues involved in the functional integrity of Escherichia coli methionine aminopeptidase. J Bacteriol 181(15):4686-4689.		
	AZ	Endo, S., et al. (2001) The additional methionine residue at the N-terminus of bacterially expressed human interleukin-2 affects the interaction between the N- and C-termini. Biochemistry, 40:914-919.		
	AAA	Fischer, W.H., and Spiess, J. (1987) Identification of a mammalian glutaminyl cyclase converting glutaminyl into pyroglutamyl peptides. Proc. Natl. Acad. Sci. U.S.A., 84:3628-3632.		
	ABB	Hirel, P.H., et al. (1989) Extent of N-terminal methionine excision from Escherichia coli proteins is governed by the side-chain length of the penultimate amino acid. Proc. Natl. Acad. Sci. U.S.A., 86:8247-8251.		
	ACC	Huang, H.C., et al. (1998) The Rana catesbeiana rcr gene encoding a cytotoxic ribonuclease. Tissue distribution, cloning, purification, cytotoxicity, and active residues for RNase activity. J. Biol. Chem., 273(11):6395-6401.		
	ADD	Hwang, D.D.W., et al. (1999) Co-expression of glutathione S-transferase with methionine aminopeptidase: a system of producing enriched N-terminal processed proteins in <i>Escherichia coli</i> . Biochem. J., 338(Pt 2):335-342.		
$\sqrt{}$	AEE	Ishitani, M., et al. (2000) SOS3 function in plant salt tolerance requires N-myristoylation and calcium binding. Plant Cell, 12:1667-1677.		
www	AFF	Leu, Y.J., et al. (2003) Residues involved in the catalysis, base specificity, and cytotoxicity of ribonuclease from Rana catesbeiana based upon mutagenesis and X-ray crystallography. J. Biol. Chem., 278(9):7300-7309.		

Examiner Signature,	Date Considered
allhin (1/1/1/2000)	2 December 2000
EXAMINER: Initials citation considered. Craw line through citation if no	t in conformance and not considered. Include copy of this form with
next communication to applicant.	

Substitute Form PTO-1449 (Modified)	U.S. Department of Commerce Patent and Trademark Office	Attorney's Docket No. 16743-004001	Application No. 10/813,549
Information Disclosure Statement by Applicant		Applicant You-Di Liao	
	eets if necessary)	Filing Date March 29, 2004	Group Art Unit . Unknown

	Other D	ocuments (include Author, Title, Date, and Place of Publication)
Examiner Initial	Desig. ID	Document
Www	AGG	Li, X., and Chang, Y.H. (1995) Amino-terminal protein processing in Saccharomyces cerevisiae is an essential function that requires two distinct methionine aminopeptidases. Proc. Natl. Acad. Sci. U.S.A., 92:12357-12361.
	АНН	Liao, Y.D., and Wang, J.J. (1994). Yolk granules are the major compartment for bullfrog (<i>Rana catesbeiana</i>) oocyte-specific ribonuclease. Eur J Biochem., 222:215-220.
	AII	Liao, Y.D., et al. (2000) Purification and cloning of cytotoxic ribonucleases from Rana catesbeiana (bullfrog). Nucleic Acids Res., 28(21):4097-4104.
	AJJ	Liao, Y.D., et al. (2003) The structural integrity exerted by N-terminal pyroglutamate is crucial for the cytotoxicity of frog ribonuclease from Rana pipiens. Nucleic Acids Res., 31(18):5247-5255.
	AKK	Lowther, W.T., et al. (1999) Escherichia coli methionine aminopeptidase: implications of crystallographic analyses of the native, mutant, and inhibited enzymes for the mechanism of catalysis. Biochemistry, 38:7678-7688.
	ALL	Lowther, W.T., and Matthews, B.W. (2000) Structure and function of the methionine aminopeptidases. Biochim. Biophys. Acta., 1477:157-167.
	AMM	Moerschell, R.P., et al. (1990) The specificities of yeast methionine aminopeptidase and acetylation of amino-terminal methionine in vivo. Processing of altered iso-1-cytochromes c created by oligonucleotide transformation. J. Biol. Chem., 265(32):19638-19643.
	ANN	Notomista, E., et al. (1999) Effective expression and purification of recombinant onconase, an antitumor protein. FEBS Lett., 463:211-215.
	A00	Prchal, J.T., et al. (1986) Hemoglobin Long Island is caused by a single mutation (adenine to cytosine) resulting in a failure to cleave amino-terminal methionine. Proc. Natl. Acad. Sci. U.S.A., 83:24-27.
	APP	Roderick, S.L., and Matthews, B.W. (1993) Structure of the cobalt-dependent methionine aminopeptidase from <i>Escherichia coli</i> : a new type of proteolytic enzyme. Biochemistry, 32:3907-3912.
	AQQ	Shapiro, R., et al. (1988) Expression of Met-(-1) angiogenin in Escherichia coli: conversion to the authentic <glu-1 175:450-461.<="" anal.="" biochem.,="" protein.="" td=""></glu-1>
	ARR	Tahirov, T.H., et al. (1998) Crystal structure of methionine aminopeptidase from hyperthermophile, <i>Pyrococcus furiosus</i> . J. Mol. Biol., 284:101-124.
\bigvee	ASS	Tobias, J.W., et al. (1991) The N-end rule in bacteria. Science, 254:1374-1377.
www	ATT	Varshavsky, A. (1996) The N-end rule: functions, mysteries, uses. Proc. Natl. Acad. Sci. U.S.A., 93:12142-12149.

Examiner Signature	Date O
Examiner digridule	Date Considered
1/1/1/May. 11) 1/Wolle	2 December 2005
EXAMINER: Initials diation considered. Draw line through citation if no	t in conformance and not considered. Include conv of this form with
next communication to applicant.	The second state of the contract of the second state of the second
	Substitute Disclosure Form (PTO-1449)

Substitute Form PTO-1449 (Modified) U.S. Department of Commerce Patent and Trademark Office		Attorney's Docket No. 16743-004001	Application No. 10/813,549	
Information Disclosure Statement by Applicant		Applicant You-Di Liao		
(Use several si	neets if necessary)	Filing Date March 29, 2004	Group Art Unit Unknown	

(Other Documents (include Author, Title, Date, and Place of Publication)			
Examiner Initial	Desig. ID	Document		
illong	AUU	Vetro, J.A., and Chang, Y.H. (2002) Yeast methionine aminopeptidase type 1 is ribosome-associated and requires its N-terminal zinc finger domain for normal function <i>in vivo</i> . J. Cell. Biochem., 85:678-688.		
lum	AVV	Walker, K.W., and Bradshaw, R.A. (1999) Yeast methionine aminopeptidase I. Alteration of substrate specificity by site-directed mutagenesis. J. Biol. Chem., 274(19):13403-13409.		

Examiner Signature	Date Considered
Millani. Merre	2 December Zert
EXAMINER: Initials citation considered. Draw line through citation if not in conformance and not considered. Include copy of this form with	
next communication to applicant.	